

Citation for published version:

Jones, E 2019, 'Towards the Marine Biorefinery: Seawater Fractionation of Macroalgae', Developing sustainable technologies in the bioeconomy and energy sectors, Bath, UK United Kingdom, 8/05/19 - 9/05/19.

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Publisher Rights
Unspecified

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Towards the Marine Biorefinery: Seawater Fractionation of Macroalgae.



UNIVERSITY OF
BATH



Centre for **Sustainable**
Chemical Technologies

E.S. Jones^a, C.J. Chuck^b

e-mail: E.S.Jones@bath.ac.uk: web: www.csct.ac.uk

^aEPSRC Centre for Doctoral Training, Centre for Sustainable Chemical Technologies, University of Bath, BA2 7AY, UK.

^bDepartment of Chemical Engineering, University of Bath, BA2 7AY


Why seawater?

- ◆ The **Marine Biorefinery** concept is built on seaweed and algae
- ◆ **Seaweed** is wild-harvest from the oceans, it's a better resource than terrestrial biomass crops in many ways¹
- ◆ Oceans are **saline**, 35 parts per thousand of salt on average
- ◆ Current Marine Biorefinery research takes place in the Lab in **Freshwater**
- ◆ Why not just do the basic chemistry **directly in seawater**?


Experimental Questions:

- ⌚ How does the presence of salt impact a simple model fractionation reaction?
- ⌚ Does the presence of salt influence the optimal reaction conditions?
- ⌚ Does the presence of salt inhibit a model fermentation organism?



Materials



Macroalgae: *Fucus serratus*²




Solvent: Freshwater to Brine³



1. Fractionation:

- ✓ Design of Experiment: Multifactorial screening experiment, varying salinity, reaction temperature, and choice of catalyst
- ✓ HPLC for sugars, gravimetry for yields. CHN on solids, TOC on Aq.





4D response plot: Ash-free solids conversion yield as a function of reaction temperature, catalyst choice, and solvent salinity. Optimal conversion yield found at high temperature, with NaOH, and seawater

- References:
1. K. A. Jung, S. R. Lim, Y. Kim and J. M. Park, *Bioresour. Technol.*, 2013, **135**, 182–190.
 2. "Fucus serratus2" by Stemonitis (https://commons.wikimedia.org/wiki/File:Fucus_serratus2.jpg) accessed 30-Apr-2019, shared under Creative Commons Attribution-Share Alike 2.5 <https://creativecommons.org/licenses/by-sa/2.5/legalcode>
 3. "Water salinity diagram" by Peter Summerlin (https://commons.wikimedia.org/wiki/File:Water_salinity_diagram.png) accessed 30-Apr-2019, shared under Creative Commons Attribution-Share Alike 2.5 <https://creativecommons.org/licenses/by-sa/2.5/legalcode>
 4. F. Abeln, J. Fan, V. L. Budarin, H. Briers, S. Parsons, M. J. Allen, D. A. Henk, J. Clark and C. J. Chuck, *Algal Res.*, 2019, **38**, 101411.
 5. S. Raikova, T. D. J. Knowles, M. J. Allen and C. J. Chuck, *ACS Sustain. Chem. Eng.*, 2019, **7**, 6769–6781.



2. Fermentation⁴:


- ✓ 96-Well Plate format for rapid screening of experimental space
- ✓ OD₆₀₀ absorption every 30 minutes for 5.5 days
- ✓ *Metschnikowia pulcherrima* (NCYC 4331) - inhibitor tolerant, catabolises oligo-saccharides, makes synthetic palm oil⁴




Rapid fermentation screening: *M. pulcherrima* is capable of growing on both acid and base hydrolysates, and at all salt concentrations. Base conditions make for simpler reactor design due to materials of construction compatibility.

3. Hydrothermal Liquefaction⁵:

- ✓ Run on Base hydrolysate solids only
- ✓ Increased solids loading to 13 % wt.



Protein Fraction: Use of Base treatment directs Nitrogen to Aq. Phase, improves C/N ratio for HTL operation



Conclusions:

- ◆ NaOH treatment improves HTL products, and no disadvantage vs. H₂SO₄ digestion
- ◆ Seawater (35ppt) vs freshwater has negligible effect on fractionation & fermentation
- ◆ HTL vs. Fermentation products can be directed by simple temperature control

EPSRC

Engineering and Physical Sciences
Research Council